arXiv:physics/0412024v2 [physics.class-ph] 20 Dec 2005

The Minkowski metric in non-inertial observer radar coordinates*

E. Minguzzi†

Departamento de Matemáticas, Universidad de Salamanca, Plaza de la Merced 1-4, E-37008 Salamanca, Spain and INFN, Piazza dei Caprettari 70, I-00186 Roma, Italy

We give a closed expression for the Minkowski $(1+1)$-dimensional metric in the radar coordinates of an arbitrary non-inertial observer O in terms of O's proper acceleration. Knowledge of the metric allows the non-inertial observer to perform experiments in spacetime without making reference to inertial frames. To clarify the relation between inertial and non-inertial observers the coordinate transformation between radar and inertial coordinates, also is given. We show that every conformally flat coordinate system can be regarded as the radar coordinate system of a suitable observer for a suitable parametrization of the observer worldline. Therefore, the coordinate transformation between arbitrarily moving observers is a conformal transformation and conformally invariant (1+1)-dimensional theories lead to the same physics for all observers, independently of their relative motion.

I. INTRODUCTION

Although the structure of special relativity is best explained using inertial frames, it is natural to ask how Minkowski spacetime would appear to a non-inertial observer. The motion of a non-inertial observer is often studied in an inertial reference frame. However, it is more convenient to construct coordinates for the non-inertial observer and to express the metric in these coordinates. In this way the non-inertial observer does not need to communicate directly with inertial observers to interpret the results of her own experiments. Two observations will guide the mathematical derivations of this work. (a) The definition of the coordinates must be operationally clear. In (1+1)-dimensions there are many possible choices, but only radar coordinates are clear operationally. Other well known coordinates, such as Fermi coordinates,[5,6,7,8] are mathematically convenient but turn out to be difficult to interpret operationally. (b) The motion of the non-inertial observer influences the expression of the metric in the non-inertial coordinates. It is convenient to express the motion of the non-inertial observer by means of her proper acceleration because this quantity is directly observable and makes no reference to external inertial frames. Following this strategy the non-inertial observer metric is expected to depend on the proper acceleration history of the non-inertial observer.

We shall be particularly concerned with what can be actually measured by the non-inertial observer, and our construction will depend on an observable definition of the coordinates (radar coordinates) and on an observable and intrinsic description of the non-inertial observer motion (the proper acceleration).

We consider a non-inertial observer O in unidirectional, but otherwise arbitrary, motion in the Minkowski two-dimensional spacetime of metric signature $(+-)$. The Minkowski metric is given by $ds^2 = dt^2 - dx^2$, where x and t are coordinates of an inertial frame K and where units have been chosen such that $c = 1$. Let $\gamma(\tau)$ be the arbitrary timelike worldline of the non-inertial observer, τ the proper time parametrization, and $u^\mu = dx^\mu/d\tau$ with $\mu = 0, 1$ the covariant velocity.

As mentioned, the non-inertial observer can construct coordinates using different methods. The transformation between inertial and non-inertial coordinates should reduce to a Lorentz transformation if the observer has vanishing acceleration. However, this requirement is too weak to determine a unique solution. There are at least three possibilities that have been considered: radar coordinates,[1,2,3,4] Fermi coordinates[5,6,7,8] and Møller coordinates.[9,10]

Even more possibilities arise if the observer is not regarded as isolated, but as part of an extended reference frame[11] where no worldline has a privileged role.[12,13] In this case the coordinate system also would depend on the rigidity assumptions that constrain the frame.[14,15,16,17]

Here we shall focus on radar coordinates. They are commonly used in the study of accelerating[1,4] and rotating systems in special relativity.[18,19] These coordinates are not related to measurements performed with standard rods and clocks. They are just convenient labels of spacetime events. However, in the neighborhood of the observer worldline, because they approximate the coordinates of a comoving inertial frame, they represent measurements performed by rods and clocks. For example, the radar distance becomes the distance that would be measured using standard rods.[2]

II. RADAR COORDINATES AND PROPER ACCELERATION

Radar coordinates are defined as follows. Let E be a spacetime event and consider the light beam L_2 emitted at E. It reaches O's worldline at the proper time $\tau_2(E)$. Consider also the light beam L_1 that was emitted at a suitable proper time $\tau_1(E)$ and reaches E. By using the radar procedure, the observer O assigns to E a time label given by $(\tau_2(E) + \tau_1(E))/2$. Note that if $E = \gamma(\tau)$, then the time label assigned to E is just τ. For this reason the time label will be denoted by the same symbol

$$\tau(E) = \frac{\tau_2(E) + \tau_1(E)}{2}. \tag{1}$$

Next O assigns to the event a radar distance given by

$$d(E) = \frac{\tau_2(E) - \tau_1(E)}{2}. \tag{2}$$

It is more convenient to define a spatial coordinate χ rather than a distance, and we define $\chi = \pm d$ with the plus sign if E stays to the right of the curve $\gamma(\tau)$ or the minus sign if E stays to the left. It also is convenient to introduce the quantities $\tau^+(E)$ and $\tau^-(E)$ (see Fig. II). These quantities coincide respectively with $\tau_2(E)$ and $\tau_1(E)$ if E stays to the right of the curve $\gamma(\tau)$, and in the opposite case, they coincide respectively with $\tau_1(E)$ and $\tau_2(E)$. We can say that τ^+ determines the intersection event between γ and a light beam moving from the right to the left, and τ^- determines the intersection event between γ and a light beam moving from the left to the right. With these definitions we have the radar coordinates



FIG. 1: Definition of τ^+ and τ^-.

$$\tau(E) = \frac{\tau^+(E) + \tau^-(E)}{2}, \tag{3}$$

$$\chi(E) = \frac{\tau^+(E) - \tau^-(E)}{2}, \tag{4}$$

where $\tau^+(E)$ and $\tau^-(E)$ are directly observable quantities. The radar coordinates are defined in the intersection between the causal future and the causal past of the curve γ. Outside this region, τ_2, or τ_1, or both, are not defined. A typical example is the constant acceleration case. Only the events in the Rindler wedge, $-x < t < x$, admit two light beams passing through them that intersect the curve γ, thus defining τ_2 and τ_1.

If we introduce the coordinates $x^\pm = t \pm x$, then the Minkowski metric takes the form

$$ds^2 = dx^+ dx^-. \tag{5}$$

Note that the events with a given value of $\tau^\pm$ lie in a plane of equation $x^\pm =$ constant. This fact will be central in what follows and justifies the introduction of τ^+ and τ^- in place of τ_2 and τ_1.

Finally, we introduce the acceleration of the non-inertial observer given by $a^\mu = \nabla_u u^\mu$, where ∇ is the Levi-Civita connection of the Minkowski metric $ds^2 = g_{\mu\nu}dx^\mu dx^\nu$, that is, if v is a vector field, $(\nabla_u v)^\alpha = v^\alpha_{\ ,\mu}u^\mu + \Gamma^\alpha_{\nu\mu}v^\nu u^\mu$, where $\Gamma^\alpha_{\nu\mu} = \frac{1}{2}g^{\alpha\beta}(g_{\beta\mu,\nu} + g_{\beta\nu,\mu} - g_{\mu\nu,\beta})$ are the Christoffel coefficients. In inertial frame coordinates $g_{\mu\nu} = \eta_{\mu\nu}$ and $\Gamma^\alpha_{\mu\nu} = 0$. Because $u \cdot u = 1$, we have $a \cdot u = 0$, and hence a is spacelike. Let e_1 be the normalized ($e_1 \cdot e_1 = -1$) spacelike vector orthogonal to u and oriented toward increasing x ($e_1[x] > 0$). Because $a \cdot u = 0$, we have

$$a^\mu(\tau) = a(\tau) e_1^\mu(\tau), \tag{6}$$

where a is the acceleration of O with respect to the local inertial frame. In other words, $-a(\tau)$ is the apparent acceleration of free particles with respect to the accelerated observer.

III. THE MINKOWSKI METRIC IN RADAR AND FERMI COORDINATES

Our goal is to prove the the following result: In the radar coordinates (τ, χ), the Minkowski metric takes the form

$$ds^2 = e^{\int_{\tau-\chi}^{\tau+\chi} a(\tau')d\tau'}(d\tau^2 - d\chi^2). \tag{7}$$

This result generalizes to arbitrary motion the constant acceleration case considered by Lass[20] for which he finds, given certain assumptions, that

$$ds^2 = e^{2a\chi}(d\tau^2 - d\chi^2). \tag{8}$$

The fact that the coordinates τ and χ are radar coordinates of the uniformly accelerated observer was pointed out later.[21] The constant acceleration case is also discussed in Refs. 15,22,23,24. It is remarkable that the problem can be solved without restricting assumptions. Note that at time τ the non-inertial observer knows the spacetime coordinates (τ, χ) of the events in the causal past of $\gamma(\tau)$. At the same time the observer can immediately calculate the metric in its causal past from the knowledge of the acceleration a as measured by a comoving accelerometer. It also is interesting that the metric at the event E is determined by the acceleration of the observer in events that are spacelike separated from E. Indeed, the integral that appears in Eq. (7) is between $\tau^-(E)$ and $\tau^+(E)$. From Eq. (7) we see that the radar coordinates are conformally related with the inertial coordinates. Thus the speed of light in radar coordinates is equal to one in both directions,[15] indeed $ds^2 = 0$ implies $|d\chi/d\tau| = 1$.

The proof of (7) goes as follows. Because the level set of the function $\tau^\pm$ coincides with that of the function $x^\pm$, there exist functions g^+ and g^- such that

$$x^+ = g^+(\tau^+), \tag{9a}$$

$$x^- = g^-(\tau^-). \tag{9b}$$

We therefore have

$$\begin{aligned} ds^2 &= {g^+}'(\tau^+){g^-}'(\tau^-)d\tau^+ d\tau^- \qquad (10a)\\ &= {g^+}'(\tau^+){g^-}'(\tau^-)(d\tau^2 - d\chi^2). \qquad (10b)\end{aligned}$$

The worldline γ satisfies $\chi = 0$, and along this worldline, $ds = d\tau$ and $\tau^+ = \tau^- = \tau$; thus ${g^+}'(\tau){g^-}'(\tau) = 1$. This argument, which also can be found in Jones,[25] allows us to conclude that there is a function $G(\tau)$ such that

$$g^\pm(\tau) = \int_0^\tau e^{\pm G(\tau')} d\tau' + C^\pm, \qquad (11)$$

where $C^\pm$ is an integration constant. The metric takes the form

$$ds^2 = e^{G(\tau+\chi) - G(\tau-\chi)}(d\tau^2 - d\chi^2). \qquad (12)$$

This metric shows that $e_1 = \partial/\partial\chi$ along the worldline $\chi = 0$.

Now we make a little digression and consider a different problem. Let $g'_{\mu\nu} = \Omega^2 g_{\mu\nu}$ be two conformally related metrics. The geodesics of $g'_{\mu\nu}$ are extremals of the length functional

$$I[\gamma] = \int_\gamma \sqrt{g'_{\mu\nu}\frac{dx^\mu}{d\lambda}\frac{dx^\nu}{d\lambda}}\, d\lambda = \int_\gamma \Omega \sqrt{g_{\mu\nu}\frac{dx^\mu}{d\lambda}\frac{dx^\nu}{d\lambda}}\, d\lambda. \qquad (13)$$

Let $u^\mu = dx^\mu/ds$ with $ds^2 = g_{\mu\nu}dx^\mu dx^\nu$. If we take the variational derivative of $I[\gamma]$, we obtain

$$g_{\mu\nu}\nabla_u u^\nu = (\delta^\alpha_\mu - u^\alpha u_\mu)\partial_\alpha \ln\Omega, \qquad (14)$$

where ∇ is the Levi-Civita connection of $g_{\mu\nu}$.

We return to our original problem, and let $g = dt^2 - dx^2$ be the Minkowski metric, $g' = d\tau^2 - d\chi^2$, and $\Omega^{-2} = e^{G(\tau+\chi) - G(\tau-\chi)}$. The worldline $\chi = 0$ is clearly a geodesic for the metric g', and therefore if u^μ and a^μ are the covariant velocity and acceleration of O, we have

$$a_\mu = -\frac{1}{2}(\delta^\alpha_\mu - u^\alpha u_\mu)\big[\partial_\alpha G(\tau+\chi)|_{\chi=0} - \partial_\alpha G(\tau-\chi)|_{\chi=0}\big]. \qquad (15)$$

The product of (15) with e_1^μ gives

$$a(\tau) = \frac{1}{2}[\partial_\chi G(\tau+\chi)|_{\chi=0} - \partial_\chi G(\tau-\chi)|_{\chi=0}] = G'(\tau), \qquad (16)$$

from which it follows that

$$G(\tau) = \int_0^\tau a(\tau')d\tau' + G(0). \qquad (17)$$

If we substitute Eq. (17) into Eq. (12), we obtain Eq. (7).

The coordinate transformation between (τ,χ) and (t,x) can be found from Eqs. (9) and (11)

$$\begin{aligned} t \;&=\; t(0) + \frac{1}{2}\sqrt{\frac{1+v(0)}{1-v(0)}}\int_0^{\tau+\chi} e^{\int_0^{\tau'} a(\tau'')d\tau''} d\tau' \\ &\quad + \frac{1}{2}\sqrt{\frac{1-v(0)}{1+v(0)}}\int_0^{\tau-\chi} e^{-\int_0^{\tau'} a(\tau'')d\tau''} d\tau', \qquad (18)\\ x \;&=\; x(0) + \frac{1}{2}\sqrt{\frac{1+v(0)}{1-v(0)}}\int_0^{\tau+\chi} e^{\int_0^{\tau'} a(\tau'')d\tau''} d\tau' \\ &\quad - \frac{1}{2}\sqrt{\frac{1-v(0)}{1+v(0)}}\int_0^{\tau-\chi} e^{-\int_0^{\tau'} a(\tau'')d\tau''} d\tau', \qquad (19)\end{aligned}$$

where $v(0)$ and and $(t(0), x(0))$ are the velocity and position of O with respect to K at $\tau = 0$. For $a = 0$ this transformation becomes a Poincaré transformation between an inertial frame moving at speed $v(0)$ and K.

It is interesting to compare these results with the analogous results obtained for the Fermi coordinates. The accelerated observer sets up Fermi coordinates (τ_f, χ_f) if two conditions are fulfilled: (1) the simultaneity slices, τ_f = constant, are lines perpendicular to the observer worldline, and (2) given the event E, $\chi_f(E)$ is, up to a sign, the length of the geodesic segment that lies in the simultaneity slice and that connects E with the observer's worldline. In the $1+1$ case, the coordinate transformation from Fermi to inertial coordinates was derived in Ref. 26 and studied in Ref. 27. The main result of Refs. 26 and 27 is quite elegant. We summarize it in the following. For a proof we refer the reader to the original work[26,27] or to the problems in Sec. VI where a more direct derivation is proposed.

Let $(T(\tau), X(\tau))$ be the trajectory of the non-inertial observer O with respect to an inertial frame of coordinates (t,x), where τ is O's proper time, and let (τ_f, χ_f) be Fermi coordinates set up by O. Then

$$t = T(\tau_f) + \chi_f \dot{X}(\tau_f), \qquad (20)$$

$$x = X(\tau_f) + \chi_f \dot{T}(\tau_f), \qquad (21)$$

and the metric takes the form

$$ds^2 = [1 + a(\tau_f)\chi_f]^2 d\tau_f^2 - d\chi_f^2, \qquad (22)$$

where $a(\tau)$ is the acceleration with respect to the local inertial frame. For a constant acceleration a, this metric is well known. It is remarkable that it changes only minimally, through the dependence of a on the proper time, when a is allowed to change.

IV. AN INVERSE RESULT

The radar coordinates of O are obtained from the proper time measurements τ^+ and τ^-. Suppose that O parametrizes the worldline γ using a new parameter $\mathcal{T} = f(\tau)$ with $f' > 0$. Moreover, suppose that O still makes use of the radar method in order to extend a coordinate system over spacetime. In place of $\tau^{\pm}$, the observer measures $\mathcal{T}^{\pm} = f(\tau^{\pm})$ and assigns to the generic event E with coordinates (τ, χ) the coordinates

$$\mathcal{T} = \frac{f(\tau+\chi) + f(\tau-\chi)}{2}, \tag{23}$$

$$\mathcal{X} = \frac{f(\tau+\chi) - f(\tau-\chi)}{2}. \tag{24}$$

Transformations of this form are called *worldline reparametrizations* independently of the physical interpretation of the coordinates. Given a worldline γ, the worldline reparametrizations form a group of coordinate transformations; in this group there is a particular choice of coordinates that corresponds to the proper time parametrization that ultimately is used in practice. In what follows all the coordinate systems in the group will be considered as radar coordinates for the worldline γ. We shall say that $(\mathcal{T}, \mathcal{X})$ are radar coordinates of the parametrized curve $\gamma(\mathcal{T})$, while (τ, χ) are radar coordinates of the parametrized curve $\gamma(\tau)$ (proper radar coordinates).

We found in Sec. III that the radar coordinates are conformally related with the inertial coordinates. Now we give a result that may be thought of as the inverse of this result. We recall that $(\mathcal{T}, \mathcal{X})$ are conformally flat coordinates if the metric (in our case the Minkowski metric) takes the form $ds^2 = \Omega(\mathcal{T}, \mathcal{X})^2(d\mathcal{T}^2 - d\mathcal{X}^2)$ for a suitable positive function Ω. Let R be the coordinate transformation from the inertial coordinates (t, x) to the conformally flat coordinates $(\mathcal{T}, \mathcal{X})$, that is, $(\mathcal{T}, \mathcal{X}) = R(t, x)$, and define the coordinates $(\mathcal{T}', \mathcal{X}') = R \circ P(t, x)$, where P is a Poincarè transformation. Then the spacetime metric in the primed coordinates has the same functional dependence as the unprimed coordinates, $ds^2 = \Omega(\mathcal{T}', \mathcal{X}')^2(d\mathcal{T}'^2 - d\mathcal{X}'^2)$. We stress that $(\mathcal{T},\ \mathcal{X})$, and $(\mathcal{T}', \mathcal{X}')$ are distinct conformally flat coordinates.

Note that if $(\mathcal{T}, \mathcal{X})$ is a conformally flat coordinate system, then $\partial_{\mathcal{T}}$ is timelike and either past-pointing or future-pointing. Analogously $\partial_{\mathcal{X}}$ is spacelike and either left-pointing or right-pointing. We restrict our attention to a conformally flat coordinate system for which $\partial_{\mathcal{T}}$ is future-pointing and $\partial_{\mathcal{X}}$ is right-pointing. Otherwise, we have only to make the replacement $\mathcal{T} \to -\mathcal{T}$ or/and $\mathcal{X} \to -\mathcal{X}$. We shall call them *oriented conformally flat coordinate systems.*

Our goal is to prove the following statement: If $(\mathcal{T}, \mathcal{X})$ are oriented conformally flat coordinates in $1+1$ Minkowski spacetime, then $(\mathcal{T}, \mathcal{X})$ are radar coordinates of the parametrized curve $\gamma(\mathcal{T})$ of the equation $\mathcal{X} = 0$. An important consequence of this result is that up to worldline reparametrizations, there is a one-to-one correspondence between oriented conformally flat coordinate systems in $1+1$ Minkowski spacetime and observers (timelike worldlines).

The proof goes as follows. Define $\mathcal{T}^{\pm} = \mathcal{T} \pm \mathcal{X}$, the level sets $\mathcal{T}^{\pm} =$ constant are null, and hence $\mathcal{T}^{\pm}$ is a invertible function of $x^{\pm}$. (We use the fact that $\partial_{\mathcal{T}}$ is future-pointing and $\partial_{\mathcal{X}}$ is right-pointing.) If we use this functional dependence in the metric, we obtain $dx^+dx^- = \Omega^2 d\mathcal{T}^+ d\mathcal{T}^-$, and it follows that $\Omega^2 = \exp[H^+(\mathcal{T}^+) + H^-(\mathcal{T}^-)]$ for suitable functions H^- and H^+. The same result could have been derived by noting that the Ricci scalar in two spacetime dimensions reads,[29] $R = -\Omega^{-2}(\partial^2_{\mathcal{T}} - \partial^2_{\mathcal{X}}) \ln \Omega^2$; because we are assuming a flat metric, $R = 0$. We define the functions

$$F = \frac{H^+ + H^-}{2}, \tag{25}$$

$$\tilde{G} = \frac{H^+ - H^-}{2}. \tag{26}$$

Then

$$ds^2 = e^{\tilde{G}(\mathcal{T}^+) - \tilde{G}(\mathcal{T}^-)} e^{F(\mathcal{T}^+) + F(\mathcal{T}^-)} d\mathcal{T}^+ d\mathcal{T}^-. \tag{27}$$

Next, introduce the coordinates (τ, χ) and the corresponding $\tau^{\pm}$ through the worldline reparametrization, Eqs. (23) and (24), or $\tau^{\pm} = f^{-1}(\mathcal{T}^{\pm})$, $f^{-1}(f(y)) = y$ with $f^{-1}(\mathcal{T}) = \int_0^{\mathcal{T}} e^{F(\mathcal{T}')} d\mathcal{T}'$. In addition, we define $G(\tau^{\pm}) = \tilde{G}(f(\tau^{\pm}))$ and obtain

$$ds^2 = e^{G(\tau^+) - G(\tau^-)} d\tau^+ d\tau^-. \tag{28}$$

Let γ be the timelike curve $\mathcal{X} = 0$. The Eq. (28) shows that τ is the proper time parametrization of γ. Indeed, the spacetime functions $\mathcal{T}^{\pm}$ coincide on γ with the curve parametrization $\mathcal{T}^{\pm}|_{\gamma} = \mathcal{T}|_{\gamma}$, and hence $\tau^+|_{\gamma} = \tau^-|_{\gamma} = \tau$ and from Eq. (28), $ds = d\tau$. According to the results of Sec. III, Eq. (28) shows that the acceleration of γ is $a(\tau) = G'(\tau)$. The coordinates (τ, χ) are therefore the proper radar coordinates of γ, and hence $(\mathcal{T}, \mathcal{X})$ are radar coordinates with respect to the worldline parametrization $\gamma(\mathcal{T})$.

V. SUMMARY

We have given the transformation between the radar coordinates of an arbitrarily accelerating observer and the radar coordinates of an inertial observer. A closed expression for the spacetime metric in the former coordinates in terms of the acceleration of the non-inertial observer also has been given. Then we showed that every oriented conformally flat coordinate system represents, up to a worldline reparametrization, an observer. Because every observer is represented by an oriented conformally flat coordinate system, the coordinate transformation between two arbitrary observers is a conformal transformation. Therefore, a (1+1)-dimensional theory invariant under conformal transformations would give rise to the same physics for all the observers independently of their relative motion. This symmetry should be broken by a Lagrangian that defines a realistic physical theory. Otherwise, it would be impossible to measure the acceleration with an accelerometer.

These considerations are related to analogous conclusions that have been reached in higher dimensional cases. Because the conformal group has only fifteen generators for more than two dimensions, it can represent only the coordinate transformations between uniformly accelerated observers.[30,31] (This case should be compared with the two-dimensional case considered here where the conformal group has an infinite number of generators.) Because Maxwell's equations are invariant under conformal transformations, they determine a physics that would appear to be the same in uniformly accelerating and inertial frames. However, the mass term in the Lorentz force equation ultimately spoils the symmetry, leading to the possibility of determining the acceleration of a frame by looking at the motion of charged particles.[32,33]

VI. SUGGESTED PROBLEMS

Problem 1. Verify that Eqs. (18) and (19) reduce to the Lorentz transformation for $a(\tau) = 0$, $t(0) = x(0) = 0$.

Problem 2. Verify that the coordinates $(\tau_f,\ \chi_f)$, which satisfy Eqs. (20) and (21), are Fermi coordinates by proving that (a) the simultaneity slices, $\tau_f =$ constant, are lines orthogonal to the observer worldline. (Use the fact that the observer's covariant velocity $u = (\dot{T}, \dot{X})$ is orthogonal to the normalized spacelike vector $e_1 = (\dot{X}, \dot{T})$ that enters in the transformation law in vectorial form.) (b) The distance of the observer worldline $(T(\tau), X(\tau))$ from the event (t, x) along the simultaneity slice is given by χ_f (because τ is a proper time $\dot{T}^2 - \dot{X}^2 = 1$). Recall from Eq. (6) that $a^\mu = a(\tau) e_1^\mu$, where $a(\tau)$ is the acceleration of O with respect to the local inertial frame. Prove that $\dot{T}(\tau) = a(\tau)\dot{X}(\tau)$ and $\dot{X}(\tau) = a(\tau)\dot{T}(\tau)$ and use these relations to derive Eq. (22).

Acknowledgments

The author is supported by INFN, grant # 9503/02.

---

[*] The published version is: E. Minguzzi, Am. J. Phys. **73**, 1117-1121 (2005).

[†] Electronic address: minguzzi@usal.es

[1] C. E. Dolby and S. F. Gull, "On radar time and the twin 'paradox'," Am. J. Phys. **69**, 1257–1261 (2001).

[2] L. D. Landau and E. M. Lifshitz, *The Classical Theory of Fields* (Addison-Wesley, Reading, MA, 1962).

[3] R. F. Märzke and J. A. Wheeler, "Gravitation as geometry: The geometry of space-time and the geometrodynamical standard meter," in *Gravitation and Relativity*, edited by H.-Y. Chiu and W. F. Hoffmann (Benjamin, New York, 1964), p. 40.

[4] M. Pauri and M. Vallisneri, "Märzke-Wheeler coordinates for accelerated observers in special relativity," Found. Phys. Lett. **13**, 401–425 (2000).

[5] E. Fermi, "Sopra i fenomeni che avvengono in vicinanza di una linea oraria" ("On the phenomena occurring near a worldline"), Rend. Accad. Lincei **31**, 21–23, 51–52, 101–103 (1922).

[6] K. P. Marzlin, "Fermi coordinates for weak gravitational fields," Phys. Rev. D **50**, 888–891 (1994).

[7] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation* (Freeman, San Francisco, 1973).

[8] A. I. Nesterov, "Riemann normal coordinates, Fermi reference system and the geodesic deviation equation," Class. Quantum Grav. **16**, 465–467 (1999).

[9] R. W. Brehme, "A geometric representation of Galilean and Lorentz transformations," Am. J. Phys. **30**, 489–496 (1962).

[10] F. W. Sears and R. Brehme, *Introduction to the Theory of Relativity* (Addison-Wesley, Reading, MA, 1968).

[11] An extended frame is defined by a congruence of timelike curves, while an observer in 1+1 Minkowski spacetime is defined by a worldline.

[12] S. Manoff, "Frames of reference in spaces with affine connections and metrics," Class. Quantum Grav. **18**, 1111–1125 (2001).

[13] E. Minguzzi, "Simultaneity and generalized connections in general relativity," Class. Quantum Grav. **20**, 2443–2456 (2003).

[14] L. Bel and J. Llosa, "Spatially affine motions in relativity," Class. Quantum Grav. **12**, 1949–1967 (1995).

[15] J. E. Romain, "Time measurements in accelerated frames of reference," Rev. Mod. Phys. **35**, 376–389 (1963).

[16] J. E. Romain, "Rigidity criteria for one-dimensional translations," Nuovo Cimento **31**, 1060–1069 (1964).

[17] N. Rosen, "Notes on rotation and rigid bodies in relativity theory," Phys. Rev. **71**, 54–58 (1947).

[18] D. G. Ashworth and R. C. Jennison, "Surveying in rotating systems," J. Phys. A **9**, 35–43 (1976).

[19] P. A. Davies, "Measurements in rotating systems," J. Phys. A **9**, 951–960 (1976).

[20] H. Lass, "Accelerating frames of reference and the clock paradox," Am. J. Phys. **31**, 274–276 (1963).

[21] E. A. Desloge and R. J. Philpott, "Uniformly accelerated reference frames in special relativity," Am. J. Phys. **55**, 252–261 (1987).

[22] L. M. Marsh, "Relativistic accelerated systems," Am. J. Phys. **33**, 934–938 (1965).

[23] J. E. Romain, "Remarks on a coordinate transformation to an accelerated frame of reference," Am. J. Phys. **32**, 279–285 (1964).

[24] D. Tinbrook, "General coordinatisations of the flat spacetime of constant proper-acceleration," Aust. J. Phys. **50**, 851–868 (1997).

[25] R. T. Jones, "Conformal coordinates associated with uniformly accelerated motion," Am. J. Phys. **24**, 124–125 (1961).

[26] M. Born and W. Biem, "Zum Uhrenparadoxon" ("On the clock paradox"), Nederl. Akad. Wetensch. Proc. Ser. B **61**, 110–120 (1958).

[27] J. Crampin, W. McCrea, F. McNally, and D. McNally, "A

class of transformations in special relativity," Proc. Roy. Soc. London. Ser. A **252**, 156–176 (1959).

[28] C. Møller, *The Theory of Relativity* (Clarendon Press, Oxford, 1962), Sec. 8.15.

[29] R. Wald, *General Relativity* (University of Chicago Press, Chicago, 1984).

[30] E. L. Hill, "On accelerated coordinate systems in classical and relativistic mechanics," Phys. Rev. **67**, 358–363 (1945).

[31] E. L. Hill, "On the kinematics of uniformly accelerated motions and classical electromagnetic theory," Phys. Rev. **72**, 143–149 (1947).

[32] T. Fulton, F. Rohrlich, and L. Witten, "Conformal invariance in physics," Rev. Mod. Phys. **34**, 442–457 (1962).

[33] T. Fulton, F. Rohrlich, and L. Witten, "Physical consequences of a coordinate transformation to a uniformly accelerating frame," Nuovo Cimento **26**, 652–671 (1962).